

Asymptotics and Hamiltonians in a First order formalism

Abhay Ashtekar^{1,*}, Jonathan Engle^{1,2,†} and David Sloan^{1‡}

¹*Institute for Gravitation and the Cosmos,
Penn State, University Park, PA 16802, U.S.A.*

²*Albert Einstein Institute, Am Mühlenberg 1, 14776 Golm, Germany*

We consider 4-dimensional space-times which are asymptotically flat at spatial infinity and show that, in the first order framework, action principle for general relativity is well-defined *without the need of infinite counter terms*. It naturally leads to a covariant phase space in which the Hamiltonians generating asymptotic symmetries provide the total energy-momentum and angular momentum of the space-time. We address the subtle but important problems that arise because of logarithmic translations and super-translations both in the Lagrangian and Hamiltonian frameworks. As a forthcoming paper will show, the treatment of higher dimensions is considerably simpler. Our first order framework also suggests a new direction for generalizing the spectral action of non-commutative geometry.

PACS numbers: 04.20.Cv, 04.20.Ha, 04.20.Fy

I. INTRODUCTION

In most field theories the action depends only on fundamental fields and their first derivatives. By contrast, the Einstein-Hilbert action of general relativity depends also on the second derivative of the fundamental field, the space-time metric g . As a consequence, stationary points of this action do not yield Einstein's equations unless both the metric and its first derivatives are kept fixed at the boundary; strictly we do not have a well-defined variational principle. To remedy this situation, Gibbons and Hawking [1, 2] proposed that we add a surface term to the Einstein Hilbert action. We are then led to

$$S_{\text{EH+GH}}(g) = \frac{1}{2\kappa} \left(\int_{\mathcal{M}} R d^4V + 2 \int_{\partial\mathcal{M}} K d^3V + C \right). \quad (1.1)$$

Here $\kappa = 8\pi G$, \mathcal{M} is a 4-manifold representing an appropriate portion of space-time, $\partial\mathcal{M}$ its boundary, R the Ricci scalar of the metric g , K the trace of the extrinsic curvature of $\partial\mathcal{M}$, and C is an arbitrary function of the metric h induced on $\partial\mathcal{M}$ by g .

Let us restrict ourselves to cases where g has signature $- , + , + , +$ and is smooth and globally hyperbolic. We will let \mathcal{M} be the space-time region bounded between two Cauchy surfaces. If \mathcal{M} is spatially compact, by setting $C = 0$, we obtain a well-defined variational principle and a finite on-shell action. However, in the asymptotically flat case, it is well-known that this strategy has some important limitations (see e.g., [3]). In particular, the action is

*Electronic address: ashtekar@gravity.psu.edu

†Electronic address: engle@cpt.univ-mrs.fr

‡Electronic address: sloan@gravity.psu.edu

typically infinite even ‘on-shell’, and indeed even when g is the Minkowski metric. To remedy this problem, Gibbons and Hawking [1, 2] proposed an infinite subtraction: Carry out an isometric embedding of $(\partial\mathcal{M}, h)$ in Minkowski space, calculate the trace K_o of the extrinsic curvature of $\partial\mathcal{M}$ defined by the Minkowski metric, and set $C = -K_o$. Can this procedure be carried out for generic asymptotically flat Lorentzian metrics g ? Now, there is a theorem due to Weyl which guarantees that any 2-manifold with a metric of positive scalar curvature can be isometrically embedded in the Euclidean 3-space. One may imagine that this result could be extended to higher dimensional Minkowski spaces. However, this expectation is not borne out. For, in a d dimensional space-time the metric h on the boundary $\partial\mathcal{M}$ has $d(d-1)/2 - (d-1)$ degrees of freedom (after removing the diffeomorphism gauge) while the choice of embedding provides a freedom worth only one function on $\partial\mathcal{M}$. Thus, even at this heuristic level, if $d \geq 4$ the freedom is not sufficient whence this infinite subtraction procedure will not work for generic metrics g .

Over the last few years, a new set of proposals for infinite counter terms C have appeared in the literature, motivated in part by earlier work of Brown and York [4]. In particular, Kraus, Larsen, Siebelink [5] have constructed a counter-term using a (non-polynomial) function of the Ricci curvature of the boundary. Mann and Marolf [3] have introduced a counter-term which is closer to the spirit of the Gibbons-Hawking proposal. They replace K_o with the trace of a tensor field \hat{K}_{ab} which generalizes the extrinsic curvature K_{ab}^o of $\partial\mathcal{M}$ with respect to the Minkowski metric, used by Gibbons and Hawking, to situations in which the boundary may not be isometrically embeddable in Minkowski space. The Mann-Marolf procedure is motivated by the form of the Gauss Codazzi equations and is carried out entirely in the physical space-time, without recourse to any embedding. Not only do these improved actions S_{imp} lead to well-defined action principles, but they also overcome another limitation of the original proposal: Now $\delta S_{\text{imp}} = 0$ at asymptotically flat solutions for *all* permissible variations δ .

Since we are dealing just with *classical* field theories where all fields are smooth, one might wonder if there is a way to avoid infinite subtractions altogether and construct an action principle which is manifestly finite from the beginning. The first goal of this paper is to show in some detail that this is indeed possible if one uses a first order framework based on orthonormal tetrads and Lorentz connections.¹ It is worth noting that this framework constitutes the starting point of Hamiltonian loop quantum gravity as well as spin-foams. The second goal of the paper is to use this action to construct a covariant Hamiltonian framework by keeping careful track of boundary conditions.² It is well-known that the asymptotic structure is quite subtle in 4-dimensions because of the presence of logarithmic translations and super-translations [10, 11]. These subtleties lead to considerable complications in the definition of asymptotic symmetries and conserved quantities. The ensuing difficulties were overcome in discussions of conserved quantities based on field equations quite sometime ago [10–14]. However, in much of the older literature on the derivation of Hamiltonians from action, these subtleties were generally overlooked (see, e.g., [15–17]). We will fill this gap starting with the first order action principle, thus providing a self-contained Lagrangian

¹ This fact was used in [6, 7] where, however, attention was focused on the inner, isolated horizon boundary rather than at infinity.

² For a complementary treatment in the canonical framework, see [8, 9]. However, that analysis is based on self-dual, rather than real Lorentz-connections and did not address issues involving the action principle discussed above.

and Hamiltonian description without the need of any infinite counter-terms. The first order framework also has the technical advantage that calculations can be carried out using exterior calculus and tend to be significantly simpler.

The paper is organized as follows. In section II we recall the asymptotic structure at spatial infinity in 4 dimensions and in section III we introduce the Lagrangian and Hamiltonian framework using the first order framework. A careful choice of boundary conditions eliminates the logarithmic and super translations and reduces the asymptotic symmetry group to Poincaré. In section IV we calculate the Hamiltonians generating these Poincaré symmetries and in section V we discuss their properties. In particular, although they are now defined using entirely different techniques these expressions of Poincaré-momentum agree with the earlier ones [10, 12–14] obtained using field equations. In this respect results of section V are similar to those obtained by Mann, Marolf and Virmani [3, 18] using their infinite counter-term subtraction in the second order framework. Section VI summarizes the results and discusses their relevance to recent developments concerning the spectral action used in non-commutative geometry [19, 20].

II. ASYMPTOTIC STRUCTURE: SUBTLETIES IN 4 DIMENSIONS

In this section we will introduce some notation, specify our boundary conditions at spatial infinity and discuss some important complications that are peculiar to 4 dimensions.

Let η be a Minkowski metric on \mathbb{R}^4 . Since we are primarily interested in spatial infinity, we will focus on the region \mathcal{R} which lies outside the light cone of some point p in the interior. On \mathcal{R} the 4-dimensional radial coordinate ρ is given by $\rho^2 = \eta_{ab}x^ax^b$ where x^a are the Cartesian coordinates of η with p as origin. Functions f of interest will admit a power series expansion of the type

$$f(\rho, \Phi) = \sum_{n=0}^m \frac{f_n(\Phi)}{\rho^n} + o(\rho^{-m}) \quad (2.1)$$

where $\Phi = (\theta, \phi, \chi)$ are the standard angles on hyperboloids \mathcal{H} defined by $\rho = \text{const}$ and where the remainder $o(\rho^{-m})$ has the property that $\lim_{\rho \rightarrow \infty} \rho^m o(\rho^{-m}) = 0$. Such a function will be said to *admit an asymptotic expansion to order m* . A tensor field $T^{a\dots b}_{c\dots d}$ will be said to admit an asymptotic expansion to order m if all its components in the *Cartesian* chart x^a do so. Derivatives $\partial T^{a\dots b}_{c\dots d}/\partial x^e$ of such a tensor field with respect to x^e will be assumed to admit an asymptotic expansion to order $m+1$. Finally, note that the remainder in (2.1) can contain terms of the type $(h(\Phi) \ln \rho)/\rho^{m+1}$ whence $\lim_{\rho \rightarrow \infty} \rho^{m+1} o(\rho^{-n})$ need not exist.

A smooth space-time metric g on \mathcal{R} will be said to be *weakly asymptotically flat at spatial infinity* if there exists a Minkowski metric η such that outside a spatially compact world tube $g - \eta$ admits an asymptotic expansion to order 1 and $\lim_{\rho \rightarrow \infty} (g - \eta) = 0$. This condition implies that in the (ρ, Φ) chart associated with η , the line element of g can be expanded as follows:

$$g_{ab}dx^a dx^b = \left(1 + \frac{2\sigma(\Phi)}{\rho}\right) d\rho^2 + 2\rho \frac{A_i(\Phi)}{\rho} d\rho d\Phi^i + \rho^2 \left(h_{ij} + \frac{h_{ij}}{\rho}\right) d\Phi^i d\Phi^j + o(\rho^{-1}) \quad (2.2)$$

where h_{ij} is the metric on the unit (time-like) hyperboloid in Minkowski space. Given such an expansion, it is tempting to think of η as a reference metric, consider its Poincaré group as the

asymptotic symmetry group of g and define Poincaré momentum —i.e., energy-momentum and relativistic angular momentum— using Hamiltonians generating these transformations on the appropriate phase space. Indeed, this is often done in the literature. However, the problem is that the metric η is not uniquely singled out by the physical metric g . For, if g admits an asymptotic expansion (2.2) with respect to one Minkowski metric η , it also admits this expansion with respect to another Minkowski metric $\bar{\eta}$ if $\eta - \bar{\eta}$ admits an asymptotic expansion to order 1 and $\lim_{\rho \rightarrow \infty} (\eta - \bar{\eta}) = 0$. Such Minkowskian metrics will be said to be *compatible*. One might first think that the Poincaré groups of compatible metric would agree asymptotically. Unfortunately, as we now show, this is not the case in 4 dimensions.

Set

$$\bar{x}^a = L_a{}^b x^b + \ln \rho C^a + T^a + S^a(\Phi) + o(\rho^0) \quad (2.3)$$

where $L_a{}^b$ is a Lorentz transformation (i.e. $L_a{}^b L_c{}^d \eta_{bd} = \eta_{ac}$), and T^a, C^a are constant vector fields with respect to η . Let $\bar{\eta}^{ab}$ be the Minkowski metric for which \bar{x}^a is the Cartesian chart. Then it is easy to check that η and $\bar{\eta}$ are compatible. If C^a and $S^a(\Phi)$ were set to zero, \bar{x}^a would be just another Cartesian chart for η to leading order whence Poincaré groups of $\bar{\eta}$ and η would have agreed asymptotically. The term $C^a \ln \rho$ defines a *logarithmic translation* while the angle dependent translation $S^a(\phi)$ is called a *super-translation*. If C^a and/or $S^a(\Phi)$ are non-zero, the Poincaré groups of η and $\bar{\eta}$ fail to agree even at infinity. Since one can define Poincaré momentum using any of them, there is considerable ambiguity in the values of conserved quantities. The ambiguity introduced by logarithmic translations was first pointed out by Bergmann [21] while that associated with super-translations was first observed at null infinity by Bondi and Sachs [22] and later at spatial infinity in [10]. These complications arise because, for physical reasons, we can only ask that $g - \eta$ should fall off as (ρ^{-1}) in 4 dimensions. Had we demanded that they fall off as (ρ^{-2}) , we would have been forced to set C^a and $S^a(\Phi)$ to zero and the ambiguities would have disappeared. But of course, because of the positive energy theorem, in this case the only solution satisfying our putative boundary conditions would have been Minkowski space! In d space-time dimensions, the physically correct condition is that $g - \eta$ should fall off as $(\rho^{-(d-3)})$, whence there is no freedom to perform super-translations or logarithmic translations if $d > 4$.

The challenge in 4-dimensions, then, is to strengthen the boundary conditions so that this freedom is eliminated without excessively limiting permissible space-time geometries. Let us begin with the logarithmic translations. It is straightforward to verify that under $x^a \rightarrow \bar{x}^a = \ln \rho C^a$, we are led to a Minkowski metric $\bar{\eta}$ and hence the ‘barred’-version of the expansion (2.2) in which

$$\bar{\sigma}(\Phi) = \sigma(\Phi) + C^a \hat{x}_a \quad \text{where} \quad \hat{x}_a = \frac{\eta_{ab} x^b}{\rho}. \quad (2.4)$$

Note that for any non-zero C^a , $C^a \hat{x}_a$ is function on the hyperboloids \mathcal{H} which is *odd under the reflection* $\hat{x}^a \rightarrow -\hat{x}^a$. Therefore, one can eliminate the freedom to perform the logarithmic translations *by simply demanding that in the asymptotic expansion (2.2), $\sigma(\Phi)$ be even under these reflections* [11]. How stringent is this requirement on σ ? Analysis of the asymptotic structure at spatial infinity at i^o ties it to the asymptotic behavior of the ‘electric’ part $E_{ab} = C_{abcd} \hat{x}^c \hat{x}^d$ of the Weyl tensor. The leading asymptotic part of E_{ab} is a tensor field tangential to \mathcal{H} given by

$${}^3E_{ab}(\Phi) := \lim_{\rho \rightarrow \infty} \rho^3 E_{ab}. \quad (2.5)$$

It admits a scalar potential which is precisely σ : ${}^3E_{ab} = -(D_a D_b \sigma + \sigma h_{ab})$, where D is the derivative operator of (\mathcal{H}, h_{ab}) [12–14]. Now one can show that if ${}^3E_{ab}$ is reflection-symmetric,

one can always choose a flat metric η so that σ is also reflection symmetric. Thus, to eliminate logarithmic translations one only needs to ask that ${}^3E_{ab}$ be reflection symmetric and then restrict oneself to asymptotic expansions (2.2) in which σ is also reflection symmetric. Finally, the symmetry condition on ${}^3E_{ab}$ is not excessively restrictive. For example, it is satisfied if g is asymptotically a finite superposition of Schwarzschild metrics with (possibly) distinct asymptotic time translations [23].

Next, let us consider super-translations. We first note a result due to Beig and Schmidt [13]: Given a metric g that admits an expansion (2.2), one can always find another Minkowski metric (which, for simplicity, we will denote by η again) such that the off-diagonal terms vanish to leading order. Thus, in the (ρ, Φ) chart of the new Minkowski metric, we have:

$$ds^2 = (1 + \frac{2\sigma}{\rho})d\rho^2 + \rho^2(h_{ij} + \frac{{}^1h_{ij}}{\rho})d\Phi^i d\Phi^j + o(\rho^{-1}). \quad (2.6)$$

where, by our assumption above, $\sigma(\Phi)$ is reflection-symmetric on \mathcal{H} . For this metric g one can calculate the asymptotic Weyl curvature which can be decomposed into an electric and a magnetic parts. The leading order magnetic part is a tensor field tangential to \mathcal{H} , given by:

$${}^3B_{ab} = \lim_{\rho \rightarrow \infty} {}^*C_{acbd}\hat{x}^c\hat{x}^d. \quad (2.7)$$

${}^3B_{ab}$ admits a natural tensor potential $S_{ab} := {}^1h_{ab} + 2\sigma h_{ab}$ on \mathcal{H} : $B_{ab} = \epsilon_a{}^{mn}D_m S_{nb}$ where ϵ and D are the alternating tensor and the derivative operator on the hyperboloid \mathcal{H} compatible with the metric h_{ab} , and indices are raised and lowered by h_{ab} . Under super-translations, $x^a \rightarrow \bar{x}^a = x^a + S^a(\Phi)$, the field ${}^3B_{ab}$ itself is left invariant while the potential transforms non-trivially:

$$\bar{\sigma} = \sigma, \quad \bar{{}^1h}_{ab} = {}^1h_{ab} - 2D_a D_b f - 2f h_{ab}, \quad \text{so that} \quad \bar{S}_{ab} = S_{ab} - 2D_a D_b f - 2f h_{ab}, \quad (2.8)$$

where the function f on \mathcal{H} is given by $f = f^a \hat{x}_a$. Finally, one can show that if ${}^3B_{ab}$ vanishes, then this transformation property of S_{ab} is sufficient to ensure that (via a suitable super-translation) we can always choose η such that $S_{ab} = 0$, i.e., ${}^1h_{ab} = -2\sigma h_{ab}$ [10, 12–14]. The transformation property (2.8) immediately implies that this exhausts the super-translation freedom. The condition ${}^3B_{ab} = 0$ is not excessively stringent. For example, it is automatically satisfied if g is stationary or axi-symmetric [24].

Let us summarize. If the asymptotic Weyl curvature of a weakly asymptotically flat space-time (\mathcal{M}, g_{ab}) is such that its leading order term is *purely electric* (i.e. if ${}^3B_{ab} = 0$) and *reflection symmetric* on \mathcal{H} , then one can eliminate both super-translations and logarithmic translations by strengthening the boundary conditions in a natural fashion by requiring in (2.6) that σ be reflection symmetric and ${}^1h_{ab} = -2\sigma h_{ab}$. Thus, the metrics g satisfying this additional conditions can be expanded as

$$ds^2 = (1 + \frac{2\sigma}{\rho})d\rho^2 + (1 - \frac{2\sigma}{\rho})\rho^2 h_{ij} d\Phi^i d\Phi^j + o(\rho^{-1}), \quad (2.9)$$

where σ is reflection symmetric. Such space-times will be said to be *asymptotically flat at spatial infinity*. If a given physical metric g satisfies this condition with respect to two distinct Minkowski metrics η and $\bar{\eta}$, then their isometry groups agree to leading order: If K^a is a Killing vector of η , there exists a Killing vector \bar{K}^a of $\bar{\eta}$ such that $\lim_{\rho \rightarrow \infty} K^a - \bar{K}^a = 0$. Hamiltonians generating these asymptotic symmetries provide unambiguous definitions of

conserved quantities at spatial infinity. We will see that the additional conditions on σ and ${}^1h_{ab}$ are essential for surface integral expressions of the Lorentz angular momentum to be well-defined.

Thus, although the asymptotic structure is quite subtle in 4 dimensions, one can strengthen the ‘obvious’ boundary conditions to eliminate the logarithmic translations and super-translations and yet admit a very large class of physically interesting examples. (For a treatment of these issues in a canonical framework, see [25].)

III. ACTION AND THE COVARIANT PHASE SPACE

We can now construct the Lagrangian and Hamiltonian descriptions *in the first order framework*. Our basic gravitational variables will be co-triads e_a^I and Lorentz connections A_a^{IJ} on space-time \mathcal{M} . Co-tetrads e are ‘square-roots’ of metrics and the transition from metrics to tetrads is motivated by the fact that tetrads are essential if one is to introduce spinorial matter. e_a^I is an isomorphism between the tangent space $T_p(\mathcal{M})$ at any point p and a fixed internal vector space V equipped with a metric η_{IJ} with Lorentzian signature. The internal indices can be freely lowered and raised using this fiducial η_{IJ} and its inverse η^{IJ} . Each co-tetrad defines a space-time metric by $g_{ab} := e_a^I e_b^J \eta_{IJ}$ which also has signature $(-+++)$. Then the co-triad e is automatically orthonormal with respect to g . Since the connection 1-forms A take values in the Lorentz Lie algebra, $A_a^{IJ} = -A_a^{JI}$. The connection acts only on internal indices and defines a derivative operator

$$D_a k_I := \partial_a k_I + A_{aI}{}^J k_J,$$

where ∂ is a fiducial derivative operator which, as usual, will be chosen to be flat and torsion free. As fundamental fields, e and A are independent. However, the equation of motion of A implies that A is compatible with e , i.e., is fully determined by e . Therefore, boundary conditions on A are motivated by those on e . These in turn are dictated by our discussion of asymptotics in section II.

In the Lagrangian and Hamiltonian frameworks we have to first introduce the precise space of dynamical fields of interest. Let us fix, once and for all, a co-frame ${}^0e_a^I$ such that $g_{ab}^0 = \eta_{IJ} {}^0e_a^I {}^0e_b^J$ is flat. The derivative operator defined by ${}^0e_a^I$ will be denoted by ∂_a ; $\partial_a {}^0e_b^I = 0$. The cartesian coordinates x^a of g_{ab}^0 and the associated radial-hyperboloid coordinates (ρ, Φ^i) will be used in asymptotic expansions. Discussion of section II suggests that the co-triads e_a^I should admit an asymptotic expansion of order 1. This suffices to obtain well-defined 4-momentum. However, detailed analysis shows that to define the Lorentz angular momentum one needs e_a^I to admit an expansion to order 2 (see Sec. IV). Therefore, we will assume that e_a^I can be expanded as:

$$e = {}^0e(\Phi) + \frac{{}^1e(\Phi)}{\rho} + \frac{{}^2e(\Phi)}{\rho^2} + o(\rho^{-2}) \quad (3.1)$$

where ${}^1e_a^I$ is given by

$${}^1e_a^I = \sigma(\Phi) (2\rho_a \rho^I - {}^0e_a^I) \quad (3.2)$$

with a reflection symmetric $\sigma(\Phi)$ (see (2.9). Here and in what follows

$$\rho_a = \partial_a \rho \quad \text{and} \quad \rho^I = \eta^{IJ} {}^0e_J^a \rho_a. \quad (3.3)$$

To appropriate leading orders, A_a^{IJ} can be required to be compatible with e_a^I on the time-like world-tube τ_∞ at spatial infinity which is part of the boundary $\partial\mathcal{M}$ of \mathcal{M} . This leads us to require that A_a^{IJ} is of asymptotic order 3,

$$A = {}^0A(\Phi) + \frac{{}^1A(\Phi)}{\rho} + \frac{{}^2A(\phi)}{\rho^2} + \frac{{}^3A(\Phi)}{\rho^3} + o(\rho^{-3}) \quad (3.4)$$

Compatibility of A with e and flatness of 0e enables us to set ${}^0A = {}^1A = 0$ and express 2A as

$$\begin{aligned} {}^2A_a^{IJ}(\Phi) &= 2\rho^2 \partial^{[J} (\rho^{-1} {}^1e_a^{I]}) \\ &= 2\rho (2\rho^{[I} \rho_a \partial^{J]} \sigma - {}^0e_a^{[I} \partial^{J]} \sigma - \rho^{-1} {}^0e_a^{[I} \rho^{J]} \sigma) \end{aligned} \quad (3.5)$$

(In spite of the explicit factors of ρ the right side is in fact independent of ρ because $\partial_a \sigma \sim \rho^{-1} \times$ (angular derivatives of σ).) We will not need the corresponding expression of 3A in terms of e and therefore demand compatibility between A and e only via (3.5).

A. Action Principle

Consider as before the 4-manifold \mathcal{M} bounded by space-like surfaces M_1 and M_2 . We will consider smooth histories (e, A) on M such that (e, A) are asymptotically flat in the sense specified above, and are such that M_1, M_2 are Cauchy surfaces with respect to the space-time metrics g defined by e , and the pull-back of A to M_1, M_2 is determined by the pull-back of e . The last condition is motivated by the fact that, since the compatibility between e and A is an equation of motion, boundary values where this compatibility is violated are not of interest to the variational principle. Finally it is convenient to partially fix the internal gauge on the boundaries. We will fix a constant, time-like internal vector n^I so that $\partial_a n^I = 0$ and require that the histories be such that $n^a := n^I e_I^a$ is the unit normal to M_1 and M_2 .

The first order gravitational action on these histories is given by (see e.g. [26])

$$S(e, A) = -\frac{1}{2\kappa} \int_{\mathcal{M}} \Sigma^{IJ} \wedge F_{IJ} + \frac{1}{2\kappa} \int_{\partial\mathcal{M}} \Sigma^{IJ} \wedge A_{IJ}, \quad (3.6)$$

where the 2-forms Σ^{IJ} are constructed from the co-tetrads and F is the curvature A :

$$\Sigma_{IJ} := \frac{1}{2} \epsilon_{IJKL} e^K \wedge e^L \quad \text{and} \quad F_I^J = dA_I^J + A_I^K \wedge A_K^J.$$

As in more familiar field theories, the action now depends only on the fundamental fields and their first derivatives. Although the connection A itself appears in the surface term at infinity, action is in fact gauge invariant. Indeed, it is not difficult to show that the compatibility between the pull-backs to M_1 and M_2 of e and A and the property $\partial_a n^I = 0$ implies that, on boundaries M_1 and M_2 , $\Sigma^{IJ} \wedge A_{IJ} = 2K \epsilon_{abc}$ where K is the trace of the extrinsic curvature of M_1 or M_2 (see e.g., section 2.3.1 of [27]). Thus, on M_1 and M_2 , the surface term in (3.6) is *precisely the Gibbons-Hawking surface term* with $C = 0$ in (1.1). Therefore, these surface contributions are clearly gauge invariant. This leaves us with just the surface term at the time-like cylinder τ_∞ at infinity. However, since e has to tend to the fixed co-tetrad 0e at infinity, permissible gauge transformations must tend to identity

on τ_∞ . Since the surface integral on τ_∞ involves only the pull-back of A to τ_∞ , it follows immediately that this surface integral is also gauge invariant. Note incidentally that on τ_∞ this term is *not* equal to the Gibbons-Hawking surface term (because $\partial_a \rho^I$ falls off only as $1/\rho$). Therefore, even if we were to assume compatibility between e and A everywhere and pass to a second order action, (3.6) would not reduce to the Gibbons-Hawking action with $C = 0$. It is also inequivalent to the Gibbons-Hawking prescription of setting $C = K_o$ because (3.6) is well-defined without reference to any embedding in flat space.

Our boundary conditions allow us to rewrite this action as

$$S(e, A) = \frac{1}{2\kappa} \int_M d\Sigma \wedge A - \Sigma \wedge A \wedge A \quad (3.7)$$

Boundary conditions also imply that the integrand falls off as ρ^{-4} . Since the volume element on any Cauchy slice goes as $\rho^2 \sin \theta d\rho d\theta d\phi$, the action is manifestly finite *even off shell* if the two Cauchy surfaces M_1, M_2 are asymptotically time-translated with respect to each other. Such space-times \mathcal{M} are referred to as *cylindrical slabs*. In this sub-section our discussion will be restricted to such space-times.³

On the class of histories considered, it is easy to check that the functional derivatives of the action are well defined with respect to both e and A . Variation with respect to the connection yields $D\Sigma = 0$. This condition implies that the connection D defined by A acts on internal indices in the same way as the unique torsion-free connection ∇ compatible with the co-tetrad (which satisfies $\nabla_a e_b^I = 0$). When this equation of motion is satisfied, the curvature F is related to the Riemann curvature R of ∇ by

$$F_{ab}{}^{IJ} = R_{ab}{}^{cd} e_c^I e_d^J.$$

Varying the action with respect to e_a^I and taking into account the above relation between curvatures, one obtains Einstein's equations $G_{ab} = 0$. Inclusion of matter is straightforward because the standard matter actions contain only first derivatives of fundamental fields without any surface terms and the standard fall-off conditions on matter fields imply that the matter action is finite on cylindrical slabs also off shell.

B. Covariant Phase Space

We will now let \mathcal{M} be \mathbb{R}^4 . The covariant phase space Γ will consist of smooth, asymptotically flat *solutions* (e, A) *to field equations* on \mathcal{M} . Thus, in contrast to section III A, \mathcal{M} is not restricted to be a cylindrical slab nor are the pull-backs of (e, A) fixed on any Cauchy surfaces. Our task is to use the action (3.6) to define the symplectic structure Ω on this Γ .

Following the standard procedure (see, e.g. [28]), let us perform second variations of the action to associate with each phase space point (e, A) and tangent vectors $\delta_1 \equiv (\delta_1 e, \delta_1 A)$

³ If the two Cauchy surfaces bounding \mathcal{M} are asymptotically boosted with respect to one another, \mathcal{M} is called a *boosted slab*. For a boosted slab there is no guarantee that the action would be finite *off shell*. (The situation is the same in Yang-Mills theory in Minkowski space.) However, when equations of motion are satisfied the action reduces just to the surface term which, because of our boundary conditions, is proportional to $\int_{\tau_\infty} \sigma d^3V$. The asymptotic behavior *in time* of σ [23] implies that this integral is finite. Thus the on shell result of [3, 18] is recovered in this first order framework.

and $\delta_2 \equiv (\delta_2 e, \delta_2 A)$ at that point, a 3-form J on \mathcal{M} , called the symplectic current:

$$J(\gamma; \delta_1, \delta_2) = -\frac{1}{2\kappa} [\delta_1 \Sigma^{IJ} \wedge \delta_2 A_{IJ} - \delta_2 \Sigma^{IJ} \wedge \delta_1 A_{IJ}]. \quad (3.8)$$

Using the fact that the fields (e, A) satisfy the field equations and the tangent vectors δ_1, δ_2 satisfy the linearized equations off (e, A) , one can directly verify that $J(\gamma; \delta_1, \delta_2)$ is closed as guaranteed by the general procedure involving second variations. Let us now consider a portion $\tilde{\mathcal{M}}$ of \mathcal{M} bounded by two Cauchy surfaces M_1, M_2 . These are allowed to be general Cauchy surfaces so $\tilde{\mathcal{M}}$ may in particular be a cylindrical or a boosted slab in the sense of section III A. Consider now a region $\tilde{\mathcal{R}}$ within $\tilde{\mathcal{M}}$, bounded by compact portions \tilde{M}_1, \tilde{M}_2 of M_1 and M_2 and a time-like cylinder τ joining $\partial\tilde{M}_1$ and $\partial\tilde{M}_2$. Since $dJ = 0$, integrating it over $\tilde{\mathcal{R}}$ one obtains

$$\int_{\tilde{M}_1} J + \int_{\tilde{M}_2} J + \int_{\tau} J = 0 \quad (3.9)$$

The idea is to take the limit as τ expands to the cylinder τ_∞ at infinity. Suppose the first two integrals continue to exist in this limit *and* the third integral goes to zero. Then, in the limit the sum of the first two terms would vanish and, taking into account orientation signs, we would conclude that $\int_M J$ is a 2-form on Γ which is *independent* of the choice of the Cauchy surface M . This would be the desired pre-symplectic structure. However, the issue of whether the boundary conditions ensure that the integrals over Cauchy surfaces converge and the flux across τ_∞ vanishes is somewhat delicate and often overlooked in the literature.⁴ If either of these properties failed, we would not obtain a well-defined symplectic structure on Γ .

Let us first consider the integral over the time-like boundary τ . As τ tends to τ_∞ , the integrand $J_{abc} \epsilon^{abc}$ tends to

$$\lim_{\tau \rightarrow \tau_\infty} \left(\delta_{[1} \frac{{}^1 \Sigma_{ab}}{\rho} \right) \wedge \left(\delta_{2]} \frac{{}^2 A_c}{\rho^2} \right) \epsilon^{abc} = \lim_{\tau \rightarrow \tau_\infty} \epsilon_{IJKL} {}^0 e_a^K (\delta_{[1} {}^1 e_b^L) (\delta_{2]} {}^2 A_c^{IJ}) \rho^{-3} \epsilon^{abc} \quad (3.10)$$

where ϵ^{abc} is the metric compatible 3-form on τ . Since the volume element on τ goes as ρ^3 , the integral of the symplectic flux over τ has a well-defined limit. But the key question is if the limit is *zero*. If not, there would be a leakage of the current J at spatial infinity and the symplectic structure would not be well-defined. Let us evaluate this term using the expressions (3.2) and (3.5) of ${}^1 e$ and ${}^2 A$. Then the second term in (3.10) reduces to

$$\epsilon_{IJKL} {}^0 e_a^K (2\rho_b \rho^L - {}^0 e_b^L) \delta_1 \sigma (2\rho^I \rho_c \partial^J \delta_2 \sigma - {}^0 e_c^I \partial^J \delta_2 \sigma - {}^0 e_c^I \rho^J \delta_2 \sigma) \rho^{-3} \epsilon^{abc} \quad -1 \leftrightarrow 2 \quad (3.11)$$

The term containing $\delta_1 \sigma \delta_2 \sigma$ vanishes because of anti-symmetrization while the remaining terms containing derivatives of $\delta_2 \sigma$ vanish because the normal ρ^c to τ is contracted either

⁴ Furthermore, even when such issues are discussed, one often considers only the restricted action $\Omega(\delta, \delta_V)$ of the pre-symplectic structure Ω , where one of the tangent vectors, δ_V , is associated with an asymptotic symmetry V^a on \mathcal{M} in the sense discussed in section IV because, as we will see, it is this restricted action that directly enters the discussion of conserved quantities. Typically the 3-forms $\Omega(\delta_1, \delta_V)$ on \mathcal{M} have a better asymptotic behavior than the generic 3-forms $\Omega(\delta_1, \delta_2)$. However, unless $\Omega(\delta_1, \delta_2)$ is well-defined for all δ_1, δ_2 , one does not have a coherent Hamiltonian framework and cannot start constructing conserved quantities.

with ϵ^{abc} or the derivative of σ . Thus, our boundary conditions imply that the symplectic flux across τ_∞ vanishes.

The next question is whether the integral over \tilde{M}_1 (and \tilde{M}_2) continues to be well-defined in the limit as we approach M_1 (resp. M_2). The leading term is again given by the integral of (3.10) over M_1 , the only difference being that ϵ^{abc} is now the metric compatible 3-form on M_1 . Since the volume element on M_1 goes as $\rho^2 d\rho d^2\Phi$, power counting argument says that the integral of this leading term can be logarithmically divergent. However, one can again expand out the leading term as (3.11) and show that it in fact vanishes. Since the remaining integrand falls off at least as fast as $1/\rho^4$, the integral over M converges. Thus, because of our boundary conditions, we are led to a well-defined pre-symplectic structure, i.e., a closed 2-form, on Γ

$$\Omega(\delta_1, \delta_2) = \frac{1}{2\kappa} \int_M \text{Tr} [\delta_1 \Sigma \wedge \delta_2 A - \delta_2 \Sigma \wedge \delta_1 A] , \quad (3.12)$$

where M is any Cauchy surface in \mathcal{M} and trace is taken over the internal indices. Ω is not a symplectic structure because it is degenerate. The vectors in its kernel represent infinitesimal ‘gauge transformations’. The physical phase space is obtained by quotienting Γ by gauge transformations and inherits a true symplectic structure from Ω . We will not carry out the quotient however because the calculation of Hamiltonians can be carried out directly on (Γ, Ω) .

IV. GENERATORS OF ASYMPTOTIC POINCARÉ SYMMETRIES

Let V^a be a vector field on \mathcal{M} representing an asymptotic Poincaré symmetry, a Killing vector field of one of the flat metrics η_{ab} in Γ . Then at any point (e, A) of Γ , the pair $(\mathcal{L}_V e, \mathcal{L}_V A)$ of fields satisfies the linearized field equations, whence $\delta_V := (\mathcal{L}_V e, \mathcal{L}_V A)$ is a vector field on Γ . (In the definition of the Lie-derivative, internal indices are treated as scalars; thus $\mathcal{L}_V e_a^I = V^b \partial_b e_a^I + e_b^I \partial_a V^b$.) The question is whether δ_V is a phase space symmetry, i.e., whether it satisfies $\mathcal{L}_{\delta_V} \Omega = 0$.

Consider the 1-form X_V on Γ defined by

$$X_V(\delta) = \Omega(\delta, \delta_V). \quad (4.1)$$

$\mathcal{L}_{\delta_V} \Omega = 0$ on Γ if and only if X_V is closed, i.e.,

$$\mathbb{d}X_V = 0$$

where \mathbb{d} denotes the exterior derivative on (the infinite dimensional) phase space Γ . If this is the case then, up to an additive constant, the Hamiltonian is given by

$$\mathbb{d}H_V = X_V.$$

The constant is determined by requiring that all Hamiltonians generating asymptotic symmetries at the phase space point $(e, A = 0)$ corresponding to Minkowski space-time must vanish. To calculate the right side of (4.1), it is useful to note the Cartan identities

$$\mathcal{L}_V A = V \cdot F + D(V \cdot A) \quad \text{and} \quad \mathcal{L}_V \Sigma = V \cdot D\Sigma + D(V \cdot \Sigma) - [(V \cdot A), \Sigma] \quad (4.2)$$

Using these, the field equations satisfied by (e, A) and the linearized field equations for δ , one obtains the required expression of X_V :

$$X_V(\delta) := \Omega(\delta, \delta_V) = -\frac{1}{2\kappa} \oint_{S_\infty} \text{Tr} [(V \cdot A) \delta \Sigma - (V \cdot \Sigma) \wedge \delta A] . \quad (4.3)$$

Note that the expression involves integrals *only* over the 2-sphere boundary S_∞ of the Cauchy surface M (i.e., the intersection of M with the hyperboloid \mathcal{H} at infinity); there is no volume term. This is a reflection of the fact that the theory is diffeomorphism invariant.

A. Energy-Momentum

Let us begin by setting $V^a = T^a$, an infinitesimal asymptotic translation. Since $\delta \Sigma \sim 1/\rho$, $A \sim 1/\rho^2$ and since the area element of the 2-sphere S_∞ at infinity grows as ρ^2 , the first term on the right side of (4.3) vanishes in the limit and we are left with

$$X_T(\delta) := \Omega(\delta, \delta_T) = \frac{1}{2\kappa} \oint_{S_\infty} \text{Tr} [(T \cdot \Sigma) \wedge \delta A] \quad (4.4)$$

which is manifestly well-defined. Inserting the asymptotic forms of the connection (3.5) and tetrad (3.1) we find:

$$X_T(\delta) = \frac{2}{\kappa} \oint_{S_\infty} [(\rho_a T^a) n^b \partial_b \delta \sigma + \delta \sigma (n_a T^a)] d^2 S_o \quad (4.5)$$

where n_b is the unit normal to the 2-sphere S_∞ within the hyperboloid \mathcal{H} and $d^2 S_o$ is the area element of the *unit* 2-sphere. Since the only dynamical variable in the integrand is σ , we can pull the δ out of the integral and obtain the Hamiltonian H_T generating the asymptotic translation T^a :

$$H_T = \frac{2}{\kappa} \oint_{S_\infty} [(\rho_a T^a) n^b \partial_b \sigma - \sigma (n_a T^a)] d^2 S_o \quad (4.6)$$

Had we selected a translational Killing field \bar{T}^a of another flat metric $\bar{\eta}_{ab}$ in our phase space Γ , we would have obtained the same answer because $\bar{T}^a - T^a = o(\rho^{-1})$.

Taking our transformation to be a unit time-translation which is asymptotically orthogonal to the Cauchy surface M under consideration (and hence to S_∞) we find the energy to be:

$$E = \frac{2}{\kappa} \oint_{S_\infty} \sigma d^2 S_o \quad (4.7)$$

Similarly, if T^a is a space-translation which is asymptotically tangential to M , we find

$$\vec{P} \cdot \vec{T} = \frac{2}{\kappa} \oint_{S_\infty} (\rho_a T^a) (n^b D_b \sigma) d^2 S_o \quad (4.8)$$

where D is the derivative operator on the unit hyperboloid (\mathcal{H}, h_{ab}) . Note that $\rho_a T^a$ are the ‘ $\ell = 1$ ’ spherical harmonics on S_∞ determined by translations T^a .

Thus the energy momentum is determined directly by σ . It follows from [10, 12, 13] that σ satisfies the hyperbolic equation $D^a D_a \sigma + 3\sigma = 0$ on (\mathcal{H}, h_{ab}) . Thus, its initial data consists of the pair $(\sigma, \dot{\sigma} = n^a D_a \sigma)$ on a 2-sphere cross-section of \mathcal{H} . Energy is given by the

‘ Y_{00} ’ component of the first piece of this data while the momentum by the ‘ Y_{1m} ’ components of the second piece. Finally, recall that the reflection through the origin of Minkowski space induces an isometry on the unit hyperboloid. Therefore *every* solution σ to the hyperbolic equation can be decomposed into a part $\sigma_{(E)}$ which is even under this reflection and a part $\sigma_{(O)}$ which is odd. Each satisfies the hyperbolic equation separately. Our boundary conditions require that σ be even. However it is easy to verify that, even if this condition had not been imposed, only the even part $\sigma_{(E)}$ contributes non-trivially to the expressions (4.7) and (4.8) of energy and momentum.

B. Relativistic Angular Momentum

Let us now set $V^a = L^a$, an infinitesimal asymptotic Lorentz symmetry. For definiteness, we will assume that it is a Lorentz Killing field of $\eta_{ab} := \eta_{IJ} {}^I e_a^I {}^J e_b^J$ so that it is tangential to the $\rho = \text{const}$ hyperboloids \mathcal{H} . The question is whether the vector field δ_L on Γ is Hamiltonian. Let us begin by examining the 1-form X_L on Γ . Using (4.3), we have:

$$X_L(\delta) := \Omega(\delta, \delta_L) = \frac{1}{2\kappa} \lim_{\rho \rightarrow \infty} \oint_{S_\rho} \text{Tr} [(L \cdot A) \delta \Sigma + (L \cdot \Sigma) \wedge \delta A] \quad (4.9)$$

where S_ρ is the 2-sphere intersection of the $\rho = \text{constant}$ hyperboloid \mathcal{H}_ρ with the Cauchy surface M used to evaluate the symplectic structure. Now, as ρ tends to infinity, Σ has a well-defined, non-zero limit, $A \sim \rho^{-2}$, $\delta A \sim \rho^{-2}$, $\delta \Sigma \sim \rho^{-1}$ and $L \sim \rho$. Therefore the second term in (4.9) is potentially divergent. Using the asymptotic form (3.5) of δA , it follows that the second term is proportional to

$$\lim_{\rho \rightarrow \infty} \oint_{S_\rho} (\delta \sigma L^a n_a) \rho^{-2} d^2 S \quad (4.10)$$

where n_a is the unit normal to S_ρ within the hyperboloid \mathcal{H}_ρ (or equivalently, to the Cauchy surface M) and $d^2 S$ is the volume element on S_ρ (which grows as ρ^2). If we were interested in the rotational sub-group of the Lorentz group adapted to M , the vector field L^a would be tangential to M , whence this term would vanish.⁵ But for a boost, L^a is proportional to n^a whence the potential divergence survives. Recall, however, that our boundary conditions require that σ be invariant under reflection symmetry. For a Lorentz boost, on the other hand, $L^a n_a$ is proportional to Y_{1m} , and therefore odd, whence the integral vanishes. Thus, thanks to the parity condition on σ , the 1-form X_L is well-defined on Γ .

To extract the Hamiltonian from X_L , we need to pull δ out of the integral. This is possible because the reflection symmetry again implies that the potentially divergent contribution from ${}^2 A_a^{IJ}$ vanishes. Furthermore, using the asymptotic forms (3.1) and (3.5) it follows that contributions from $(L \cdot {}^1 \Sigma) \wedge \delta {}^2 A$ and $(L \cdot {}^2 A) \wedge \delta {}^1 \Sigma$ cancel each other. Consequently,

⁵ One might first think that since the integral vanishes for all rotations, by changing the Lorentz frame defining the rotation subgroup one would be able to show that the integral vanishes also for boosts. This turns out not to be correct. To handle Lorentz boosts one needs a genuinely stronger asymptotic condition. As discussed below this is provided by the reflection symmetry of σ discussed in section II. This point is often not realized because much of the literature focuses only on rotational subgroups. See, e.g. [3, 18].

$X_L = \mathbb{d}H_L$ where the Hamiltonian generating the Lorentz transformation L^a is given by

$$H_L = -\frac{1}{2\kappa} \oint_{S_\infty} \text{Tr}(\hat{L} \cdot {}^o\Sigma) \wedge {}^3A, \quad (4.11)$$

where $\hat{L}^a = L^a/\rho$ is the Lorentz Killing field on the *unit* hyperboloid (\mathcal{H}, h_{ab}) . Note that in contrast to the energy momentum, the angular momentum is not determined by the leading order deviation of (e, A) from the ground state ($e, A = 0$) but by sub-leading terms.

V. RELATION TO THE SPI FRAMEWORK

The boundary conditions we imposed in section III to construct the Lagrangian and Hamiltonian frameworks imply that the space-time admits a conformal completion with conformal factor $\omega = \rho^{-2}$ in which spatial infinity is represented by a single point i^o . The conformally rescaled metric $\hat{g}_{ab} = \omega^2 g_{ab}$ can be shown to have the regularity needed in the so-called Spi-framework⁶ [10, 12]. These conditions in turn imply that various physical fields admit a direction dependent limit as one approaches i^o in space-like directions and can therefore be regarded as smooth fields on the unit hyperboloid \mathcal{H}_o in the tangent space at i^o . The boundary conditions we imposed have been shown to eliminate the logarithmic translations [11] and supertranslations [10, 12] also in the Spi-framework, reducing the asymptotic group to the Poincaré group. Using field equations in the physical space-time, fields on \mathcal{H}_o were shown to satisfy certain equations and these were used to define Poincaré momentum in terms of the asymptotic Weyl curvature. The asymptotic field equations made it evident that these quantities are conserved, i.e., are independent of the choice of the 2-sphere cross-section of \mathcal{H}_o used in their evaluation.

In section IV, by contrast, we were led to the expressions of Poincaré momenta using Hamiltonian considerations and our final expressions are surface integrals involving asymptotic forms of triads and connections rather than the Weyl curvature. It is natural compare the underlying assumptions and ask for the relation between these quantities and those obtained in the Spi framework.

A. Energy-momentum

In the Spi framework, the total energy momentum P_a is a 4-vector in the tangent space at i^o . Let T^a be an asymptotic translation in the physical space-time. Then it defines a vector T_o^a at i^o and corresponding component $T_o^a P_a$ of the 4-momentum is given by [10, 12, 14]

$$T_o^a p_a = -\frac{1}{\kappa} \oint_S \mathbf{E}_{ab} \hat{T}_o^b n^a d^2S. \quad (5.1)$$

Here, \mathbf{E}_{ab} is the ‘electric part’ of the asymptotic Weyl curvature, $\hat{T}_o^a = h_b^a T^b$ is the conformal Killing field on the unit hyperboloid (\mathcal{H}_o, h_{ab}) in the tangent space T_{i^o} , and S any 2-sphere cross section of \mathcal{H}_o . In terms of the physical space-time fields used in this paper, \mathbf{E}_{ab} is given by $\mathbf{E}_{ab} = {}^3E_{ab} \equiv \lim_{\rho \rightarrow \infty} \rho^3 C_{ambn} \rho^m \rho^n$. Thus, as mentioned in section II, ${}^3E_{ab}$ is the

⁶ Spi stands for spatial infinity and rhymes with scri that represents null infinity.

leading order electric part of the asymptotic Weyl tensor, where the electric and magnetic decomposition is carried out using a $\rho = \text{const}$ foliation.

To relate this P_a to that defined in section IV, let us recall that σ serves as a scalar potential of ${}^3E_{ab}$:

$${}^3E_{ab} = - (D_a D_b \sigma + \sigma h_{ab}) \quad (5.2)$$

Let us choose S to be the intersection of \mathcal{H}_o with a space-like plane in T_{i^o} and let T^a be unit and orthogonal to this plane. The question then is whether $T_o^a p_a$ equals the energy (4.7) defined in section IV using Hamiltonian considerations.

The trace-free property of ${}^3E_{ab}$ implies that σ satisfies the hyperbolic equation $D^a D_a \sigma + 3\sigma = 0$ and it is straightforward to verify that, *on the specific cross-section S of \mathcal{H}_o we chose*, $D_a \hat{T}_o^b = 0$. Therefore on this S , we have: $E_{ab} \hat{T}_o^a n^b = E_{ab} \hat{T}_o^a \hat{T}_o^b = -\hat{T}_o^a D_a \hat{T}_o^b D_b \sigma + \sigma = -(\Delta + 2)\sigma$, where Δ is the Laplacian on S . Hence, we have

$$T^a P_a = \frac{2}{\kappa} \oint_S \sigma d^2 S \quad (5.3)$$

which agrees with the expression (4.7) of energy obtained from Hamiltonian considerations.

Next, let T^a be a space-translation tangential to the 3-plane whose intersection with \mathcal{H}_o defined S . Then $\hat{T}_o^a n^b h_{ab} = 0$ and $D_a V^a = 2V^a \rho_a$ on S where ρ_a is the unit normal to \mathcal{H}_o . Using the fact that $D_a n^b$ vanishes on S we obtain,

$$T^a P_a = \frac{2}{\kappa} \oint_S (\hat{T}_o^a \rho_a) (n^b D_b \sigma) d^2 S \quad (5.4)$$

which agrees with the expression (4.8) of the 3-momentum obtained from Hamiltonian considerations.

Thus, the energy-momentum obtained using Hamiltonian considerations agrees with that obtained using just the asymptotic field equations in the i^o framework. A detailed examination shows that in both frameworks one can impose substantially weaker boundary conditions to arrive at this expression of energy-momentum. In particular, one only needs to require that e be of asymptotic order just 1 (rather than 2) and, furthermore, one can drop the requirement that the $1/\rho^3$ -part of the asymptotic Weyl curvature be pure electric and reflection symmetric, thereby allowing both logarithmic translations and super-translations.

B. Relativistic Angular Momentum

The situation with angular momentum is more subtle. To define Lorentz angular-momentum one has to get rid of the super-translation ambiguity also in the Spi framework. Indeed, the procedure we followed in section II merely mimicked the Spi strategy of requiring that the $1/\rho^3$ contribution to the magnetic part B_{ab} of the physical Weyl tensor should vanish and then setting its tensor potential to zero. However, in striking contrast to what we found in the discussion of Hamiltonians H_L in section IV B, the Spi angular momentum is well-defined without having to require that σ be reflection symmetric. Indeed, the Spi expression is constructed directly from ${}^4B_{ab}$ —the $1/\rho^4$ part of B_{ab} — and this field is insensitive to logarithmic translations [11]. Furthermore, as we will now show the final Spi-expression coincides with the expression (4.11) of H_L we found in section IV B. Why then was the parity condition essential in our derivation of (4.11)? It was necessary

because, in the Hamiltonian approach, not only should the expression of angular momentum be well defined, but it should also be the generator of asymptotic Lorentz transformations. More precisely, the parity condition is needed to show that the 1-form X_L on the phase space Γ is well-defined and exact, i.e., that the right side of (4.11) has the interpretation of Hamiltonians generating Lorentz rotations on the gravitational phase space.

In the Spi framework angular momentum is constructed from the $1/\rho^4$ part of the magnetic Weyl tensor

$${}^4B_{ab} := \lim_{\rho \rightarrow \infty} \rho^4 {}^*C_{ambn} \rho^m \rho^n \quad (5.5)$$

where, as before, ρ^m is the unit normal to the $\rho = \text{const}$ hyperboloids. Every Lorentz Killing field \hat{L}^a on the unit hyperboloid is of the form $\hat{L}^a = F_o^{ab} \rho_b$ where F_o^{ab} is a constant skew tensor in Minkowski space (\mathcal{M}, η) and therefore has a dual Lorentz Killing field defined by ${}^*\hat{L}^a = {}^*F_o^{ab} \rho_b$. The angular momentum $J_{\hat{L}}$ associated with the Lorentz Killing field L^a is defined as:

$$J_{\hat{L}} = \oint_S {}^4B^{ab} {}^*\hat{L}_a \epsilon_{bmn} dS^{mn} \quad (5.6)$$

where as before S is any 2-sphere cross-section of \mathcal{H}_o and ϵ is the volume 3-form on the unit hyperboloid (\mathcal{H}_o, h_{ab}) . To relate $J_{\hat{L}}$ with the Hamiltonian H_L of (4.11), we first write the Weyl tensor in (5.5) as $C_{abcd} = F_{ab}{}^{IJ} e_{cI} e_{dJ}$ and use the fact that $F_{ab}{}^{IJ}$ is given by $F_{ab}{}^{IJ} = 2\partial_{[a} A_{b]} + [A_a, A_b]^{IJ}$. Then, by using the asymptotic expansions (3.1) and (3.5), one obtains⁷,

$$\begin{aligned} J_{\hat{L}} &= -\frac{1}{2\kappa} \oint_S \text{Tr}(\hat{L} \cdot {}^o\Sigma) \wedge {}^3A, \\ &= H_L. \end{aligned} \quad (5.7)$$

Thus the angular momentum constructed from asymptotic field equations in the Spi framework agrees with that obtained in this paper from Hamiltonian considerations. It therefore also follows that in an asymptotically flat *axi-symmetric* space-time H_L reproduces the Komar integral if L^a is chosen to be the rotational Killing field, and in an asymptotically flat *stationary* space-time, it yields the same angular momentum dipole moment as that constructed from the stationary Killing field [24].

VI. DISCUSSION

In this paper we have shown that in the first order formalism based on co-triads and Lorentz connections, the Lagrangian and Hamiltonian frameworks can be constructed without having to introduce an infinite counter term subtraction in the action. However, since in four space-time dimensions physical metrics approach the flat metric only as $1/\rho$ at spatial infinity, the obvious boundary conditions allow one to make supertranslations and logarithmic translations. The asymptotic symmetry group is then larger than the Poincaré group. If one is interested only in energy momentum, these ambiguities can be ignored because one can still single out a well-defined 4-dimensional group of asymptotic translations. To

⁷ The calculation is significantly simplified by noting that (3.1) and (3.4) imply that the $1/\rho^3$ -part ${}^3B_{ab}$ of B_{ab} vanishes, whence one can retain just the $1/\rho^4$ terms in the expression of ${}^4B_{ab}$ in terms of A .

have a well-defined angular momentum, on the other hand, the obvious boundary conditions used in much of the older literature are too naive; they have to be carefully strengthened to reduce the asymptotic symmetry group to the Poincaré group. When this is done, the Hamiltonians generating asymptotic Poincaré transformations provide us with expressions of energy-momentum and Lorentz angular momentum. These agree with the expressions obtained in the Spi framework based of asymptotic field equations [10, 12]. Therefore they also agree with conserved quantities defined by other methods in restricted cases with exact symmetries [24].

For simplicity, in this paper we focused on vacuum Einstein's equations. However, inclusion of standard matter—in particular, scalar, Maxwell and Yang-Mills fields—with standard boundary conditions used in Minkowski space is straightforward. There are no surface terms in the action associated with matter and the expressions of the Hamiltonians generating asymptotic Poincaré transformations are formally the same as the ones we found. In particular, the Hamiltonians consist entirely of 2-sphere surface integrals at spatial infinity and their integrands do not receive any explicit contributions from matter. Matter makes its presence felt through constraint equations which, in response to matter, modify the asymptotic gravitational fields. Finally, in higher dimensions, the asymptotic structure is considerably simpler because the physical metric approaches the flat metric as $1/\rho^2$ or faster. This issue is discussed in the accompanying paper [29].

We conclude with a comment. In non-commutative geometry, in place of Riemannian geometry, one introduces a spectral triplet $(\mathcal{A}, \mathcal{H}, D)$ consisting of a non-commutative C^* algebra \mathcal{A} , a representation of it on a Hilbert space \mathcal{H} and a Dirac operator D acting on \mathcal{H} . A certain choice of the triplet is made to describe (a generalization of) the standard model of particle physics together with Einstein gravity. Rather general symmetry considerations then lead to a so-called ‘spectral action’ from which dynamics can be derived [19]. It has been known for some time that an asymptotic expansion of this action can be performed to make contact with the low energy physics and the first terms reproduce the Einstein-Hilbert action with a cosmological constant. Recently it was realized [20] that the spectral action can be naturally extended to incorporate the presence of boundaries and the asymptotic expansion of the new action produces *precisely* the Einstein Hilbert action with the Gibbons-Hawking counter term for $C = 0$ (see (1.1)). This is an exciting development. However, as the discussion of section I shows, in the asymptotically flat context this action has severe limitations and the extension of the spectral action to incorporate the boundary term [20] was motivated using precisely the Hamiltonian formulation in the asymptotically flat context. More generally, the non-commutative framework has been developed primarily for the Riemannian signature and passage to the Lorentz signature is contemplated via a Wick transform in the asymptotically flat context. Therefore asymptotic considerations of [3] and this paper are directly relevant to the spectral action approach. The natural question then is: Can the spectral action framework be further generalized so that the leading terms in the asymptotic expansion yields an action which is free from the drawbacks of (1.1)? The first order framework discussed in this paper presents a natural avenue for such a generalization. Indeed, the gravitational sector of the non-commutative geometry requires a spin-bundle—and hence a frame field e —as well a Dirac operator—i.e., a spin connection A . However, in the non-commutative framework the two are in essence compatible with one another from the beginning. The question is whether one can extend the framework so that they are independent to begin with and made compatible only by equations of motion. The spectral action in such a generalization could then descend to (3.6) upon a suitable asymp-

otic expansion. Quite apart from this specific application, such a ‘first order’ framework in non-commutative geometry appears also to lead to mathematical structures which are interesting in their own right, and could provide a technical bridge between non commutative geometry and loop quantum gravity.

Acknowledgment:

We would like to thank Don Marolf and Nigel Higson for stimulating discussions. This work was supported in part by the NSF grants PHY-0456913 and OISE-0601844, the Alexander von Humboldt Foundation and the Eberly research funds of Penn State.

-
- [1] Gibbons G and Hawking S W 1997 Action integrals and partition function in quantum gravity *Phys. Rev. D* **15** 2752
 - [2] Hawking S W 1979 The path integral approach to quantum gravity in *General Relativity, an Einstein Centenary Survey* eds Hawking S W and Israel W (Cambridge: Cambridge UP)
 - [3] Mann R B and Marolf D M 2006 Holographic renormalization of asymptotically flat space-times *Class. Quantum Grav.* **23** 29277 (*Preprint hep-th/0511096*)
 - [4] Brown J D and York J W 1993 Quasi-local energy and conserved charges derived from the gravitational action *Phys. Rev. D* **47** 1407
 - [5] Kraus P, Larsen F and Siebelink R 1999 The gravitational action in asymptotically AdS and flat space-times *Nucl. Phys. B* **563** 259 (*Preprint hep-th/9906127*)
 - [6] Ashtekar A, Fairhurst S and Krishnan B 2000 Isolated horizons: Hamiltonian evolution and the first law *Phys. Rev. D* **62** 104025 (*Preprint gr-qc/0005083*)
 - [7] Ashtekar A, Beetle C and Lewandowski J 2001 Mechanics of rotating isolated horizons *Phys. Rev. D* **64** 044016 (*Preprint gr-qc/0103026*)
 - [8] Ashtekar A 1987 A new Hamiltonian formulation of general relativity *Phys. Rev. D* **36** 1587
 - [9] Thiemann T (1995) Generalized boundary conditions for general relativity for the asymptotically flat case in terms of Ashtekar variables *Class. Quant. Grav.* **12** 181 (*Preprint gr-qc/9910008*)
 - [10] Ashtekar A and Hansen R O 1978 A unified treatment of spatial and null infinity I: Universal structure, asymptotic symmetries and conserved quantities at spatial infinity *J. Math. Phys.* **19** 1542
 - [11] Ashtekar A 1985 Logarithmic ambiguities in the description of spatial infinity *Found. Phys.* **15** 419
 - [12] Ashtekar A 1980 Asymptotic structure of the gravitational field at spatial infinity in *General Relativity and Gravitation: One Hundred Years After the Birth of Albert Einstein* ed Held A (New York: Plenum)
 - [13] Beig R and Schmidt B G 1982 Einstein’s equations near spatial infinity *Commun. Math. Phys.* **87** 65-80
 - [14] Ashtekar A and Romano J D 1992 Spatial Infinity as a Boundary of Spacetime *Class. Quant. Grav.* **9** 1069-100
 - [15] Hawking S W and Horowitz G 1996 The gravitational Hamiltonian, action, entropy and surface terms *Class. Quantum Grav.* **13** 1487 (*Preprint gr-qc/9501014*)

- [16] Arnowitt R, Deser S and Misner C M 1962 The dynamics of general relativity in *Gravitation: An Introduction to Current Research* ed Witten L (New York: John Wiley and sons)
- [17] Regge T and Teitelboim C 1974 Role of surface integrals in the Hamiltonian formulation of general relativity *Ann. Phys.* **88** 286
- [18] Mann R B, Marolf D M and Virmani A 2006 Covariant counterterms and conserved charges in asymptotically flat space-times *Class. Quant. Grav.* **23** 6357–78
- [19] Chamseddine A H and Connes A 1996 A universal action formula *Phys. Rev. Lett.* **77** 4868-71
1997 *Commun. Math. Phys.* **186** 731-750
- [20] Chamseddine A H and Connes A 2007 Quantum gravity boundary terms from spectral action *Phys. Rev. Lett.* **99** 071302
- [21] Bergmann P G 1961 Gauge invariant variables in general relativity *Phys. Rev.* **124** 274
- [22] Bondi H, Van den Burg A W K and Metzner 1962 *Proc. R. Soc. (London)* A **269** 21
- [23] Ashtekar A and Penrose R 1995 Mass positivity from focussing and structure of space-like infinity Twistor Newsletter 31 in *Further Advances in Twistor Theory* vol II eds Mason L J, Hughston L P and Kobak P Z (New York: Longman and John Wiley and Sons)
- [24] Ashtekar A and Magnon A 1979 On conserved quantities in general relativity *J. Math. Phys.* **20** 793
- [25] Beig B and O’Murchadha N 1987 Poincaré group as the symmetry group of canonical general relativity *Ann. Phys. (N.Y.)* **174** 463-498
- [26] Ashtekar A 1991 *Lectures on Non-perturbative canonical gravity* Notes prepared in collaboration with R. Tate (Singapore: World Scientific)
- [27] Ashtekar A and Lewandowski J 2004 Background independent quantum gravity: A status report *Class. Quantum Grav.* **21** R53–R152
- [28] Ashtekar A Bombelli L and Reula O 1991 The covariant phase space of asymptotically flat gravitational fields in *Analysis, Geometry and Mechanics: 200 Years after Lagrange* eds Francaviglia M and Holm D (Amsterdam: North-Holland)
- [29] Ashtekar A and Sloan D 2008 Asymptotics and Hamiltonians in a First order formalism (accompanying paper).